

# NEW SOME HADAMARD'S TYPE INEQUALITIES FOR CO-ORDINATED CONVEX FUNCTIONS

MEHMET ZEKI SARIKAYA<sup>♣</sup>, ERHAN. SET<sup>■</sup>, M. EMIN OZDEMIR<sup>♦</sup>,  
AND SEVER S. DRAGOMIR<sup>▼</sup>

ABSTRACT. In this paper, we establish new some Hermite-Hadamard's type inequalities of convex functions of 2-variables on the co-ordinates.

## 1. INTRODUCTION

Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a convex mapping defined on the interval  $I$  of real numbers and  $a, b \in I$ , with  $a < b$ . the following double inequality is well known in the literature as the Hermite-Hadamard inequality:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}.$$

Let us now consider a bidimensional interval  $\Delta =: [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b$  and  $c < d$ . A mapping  $f : \Delta \rightarrow \mathbb{R}$  is said to be convex on  $\Delta$  if the following inequality:

$$f(tx + (1-t)z, ty + (1-t)w) \leq tf(x, y) + (1-t)f(z, w)$$

holds, for all  $(x, y), (z, w) \in \Delta$  and  $t \in [0, 1]$ . A function  $f : \Delta \rightarrow \mathbb{R}$  is said to be on the co-ordinates on  $\Delta$  if the partial mappings  $f_y : [a, b] \rightarrow \mathbb{R}$ ,  $f_y(u) = f(u, y)$  and  $f_x : [c, d] \rightarrow \mathbb{R}$ ,  $f_x(v) = f(x, v)$  are convex where defined for all  $x \in [a, b]$  and  $y \in [c, d]$  (see [3]).

A formal definition for co-ordinated convex function may be stated as follows:

**Definition 1.** A function  $f : \Delta \rightarrow \mathbb{R}$  will be called co-ordinated convex on  $\Delta$ , for all  $t, s \in [0, 1]$  and  $(x, y), (u, v) \in \Delta$ , if the following inequality holds:

$$\begin{aligned} & f(tx + (1-t)y, su + (1-s)v) \\ & \leq tsf(x, u) + s(1-t)f(y, u) + t(1-s)f(x, v) + (1-t)(1-s)f(y, v). \end{aligned}$$

Clearly, every convex function is co-ordinated convex. Furthermore, there exist co-ordinated convex function which is not convex, (see, [3]). For several recent results concerning Hermite-Hadamard's inequality for some convex function on the co-ordinates on a rectangle from the plane  $\mathbb{R}^2$ , we refer the reader to ([1]-[6]).

Also, in [3], Dragomir establish the following similar inequality of Hadamard's type for co-ordinated convex mapping on a rectangle from the plane  $\mathbb{R}^2$ .

---

2000 *Mathematics Subject Classification.* 26A51, 26D15.

*Key words and phrases.* convex function, co-ordinated convex mapping, Hermite-Hadamard inequality.

<sup>\*</sup>corresponding author.

**Theorem 1.** *Suppose that  $f : \Delta \rightarrow \mathbb{R}$  is co-ordinated convex on  $\Delta$ . Then one has the inequalities:*

$$\begin{aligned}
 (1.1) \quad & f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
 & \leq \frac{1}{2} \left[ \frac{1}{b-a} \int_a^b f\left(x, \frac{c+d}{2}\right) dx + \frac{1}{d-c} \int_c^d f\left(\frac{a+b}{2}, y\right) dy \right] \\
 & \leq \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx \\
 & \leq \frac{1}{4} \left[ \frac{1}{b-a} \int_a^b f(x, c) dx + \frac{1}{b-a} \int_a^b f(x, d) dx \right. \\
 & \quad \left. + \frac{1}{d-c} \int_c^d f(a, y) dy + \frac{1}{d-c} \int_c^d f(b, y) dy \right] \\
 & \leq \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4}.
 \end{aligned}$$

The above inequalities are sharp.

The main purpose of this paper is to establish new Hadamard-type inequalities of convex functions of 2-variables on the co-ordinates.

## 2. INEQUALITIES FOR CO-ORDINATED CONVEX FUNCTIONS

**Lemma 1.** *Let  $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a partial differentiable mapping on  $\Delta := [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b$  and  $c < d$ . If  $\frac{\partial^2 f}{\partial t \partial s} \in L(\Delta)$ , then the following equality holds:*

$$\begin{aligned}
 (2.1) \quad & \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx \\
 & - \frac{1}{2} \left[ \frac{1}{b-a} \int_a^b [f(x, c) + f(x, d)] dx + \frac{1}{d-c} \int_c^d [f(a, y) + f(b, y)] dy \right] \\
 & = \frac{(b-a)(d-c)}{4} \int_0^1 \int_0^1 (1-2t)(1-2s) \frac{\partial^2 f}{\partial t \partial s}(ta + (1-t)b, sc + (1-s)d) dt ds.
 \end{aligned}$$

*Proof.* By integration by parts, we get

$$\begin{aligned}
 (2.2) \quad & \int_0^1 \int_0^1 (1-2s)(1-2t) \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sc + (1-s)d) dt ds \\
 &= \int_0^1 (1-2s) \left\{ (1-2t) \frac{1}{a-b} \frac{\partial f}{\partial s} (ta + (1-t)b, sc + (1-s)d) \Big|_0^1 \right. \\
 &\quad \left. + \frac{2}{a-b} \int_0^1 \frac{\partial f}{\partial s} (ta + (1-t)b, sc + (1-s)d) dt \right\} ds \\
 &= \int_0^1 (1-2s) \left\{ -\frac{1}{a-b} \frac{\partial f}{\partial s} (a, sc + (1-s)d) - \frac{1}{a-b} \frac{\partial f}{\partial s} (b, sc + (1-s)d) \right. \\
 &\quad \left. + \frac{2}{a-b} \int_0^1 \frac{\partial f}{\partial s} (ta + (1-t)b, sc + (1-s)d) dt \right\} ds \\
 &= \frac{1}{b-a} \left\{ \int_0^1 (1-2s) \left( \frac{\partial f}{\partial s} (a, sc + (1-s)d) + \frac{\partial f}{\partial s} (b, sc + (1-s)d) \right) ds \right. \\
 &\quad \left. - 2 \int_0^1 \int_0^1 (1-2s) \frac{\partial f}{\partial s} (ta + (1-t)b, sc + (1-s)d) dt ds \right\}.
 \end{aligned}$$

Thus, again by integration by parts in the right hand side of (2.2), it follows that

$$\begin{aligned}
 (2.3) \quad & \int_0^1 (1-2s) \left( \frac{\partial f}{\partial s} (a, sc + (1-s)d) + \frac{\partial f}{\partial s} (b, sc + (1-s)d) \right) ds \\
 &- 2 \int_0^1 \int_0^1 (1-2s) \frac{\partial f}{\partial s} (ta + (1-t)b, sc + (1-s)d) dt ds \\
 &= (1-2s) \frac{(f(a, sc + (1-s)d) + f(b, sc + (1-s)d))}{c-d} \Big|_0^1 \\
 &+ \frac{2}{c-d} \int_0^1 (f(a, sc + (1-s)d) + f(b, sc + (1-s)d)) ds \\
 &- 2 \int_0^1 \left\{ (1-2s) \frac{f(ta + (1-t)b, sc + (1-s)d)}{c-d} \Big|_0^1 \right. \\
 &\quad \left. + \frac{2}{c-d} \int_0^1 f(ta + (1-t)b, sc + (1-s)d) ds \right\} dt
 \end{aligned}$$

$$\begin{aligned}
 &= -\frac{f(a, c) + f(b, c)}{c - d} - \frac{f(a, d) + f(b, d)}{c - d} \\
 &\quad + \frac{2}{c - d} \int_0^1 (f(a, sc + (1 - s)d) + f(b, sc + (1 - s)d)) ds \\
 &\quad - 2 \int_0^1 \left\{ -\frac{f(ta + (1 - t)b, c)}{c - d} - \frac{f(ta + (1 - t)b, d)}{c - d} \right. \\
 &\quad \left. + \frac{2}{c - d} \int_0^1 f(ta + (1 - t)b, sc + (1 - s)d) ds \right\} dt \\
 &= \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{(d - c)} \\
 &\quad + \frac{4}{(d - c)} \int_0^1 \int_0^1 f(ta + (1 - t)b, sc + (1 - s)d) ds dt \\
 &\quad - \frac{2}{(d - c)} \left\{ \int_0^1 (f(a, sc + (1 - s)d) + f(b, sc + (1 - s)d)) ds \right. \\
 &\quad \left. + \int_0^1 (f(ta + (1 - t)b, c) + f(ta + (1 - t)b, d)) dt \right\}.
 \end{aligned}$$

Writing (2.3) in (2.2), using the change of the variable  $x = ta + (1 - t)b$  and  $y = sc + (1 - s)d$  for  $t, s \in [0, 1]^2$ , and multiplying the both sides by  $\frac{(b-a)(d-c)}{4}$ , we obtain (2.1), which completes the proof.  $\square$

**Theorem 2.** Let  $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a partial differentiable mapping on  $\Delta := [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b$  and  $c < d$ . If  $\left| \frac{\partial^2 f}{\partial t \partial s} \right|$  is a convex function on the co-ordinates on  $\Delta$ , then one has the inequalities:

$$\begin{aligned}
 (2.4) \quad &\left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 &\quad \left. + \frac{1}{(b - a)(d - c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\
 &\leq \frac{(b - a)(d - c)}{16} \left( \frac{\left| \frac{\partial^2 f}{\partial s \partial t}(a, c) \right| + \left| \frac{\partial^2 f}{\partial s \partial t}(a, d) \right| + \left| \frac{\partial^2 f}{\partial s \partial t}(b, c) \right| + \left| \frac{\partial^2 f}{\partial s \partial t}(b, d) \right|}{4} \right)
 \end{aligned}$$

where

$$A = \frac{1}{2} \left[ \frac{1}{b - a} \int_a^b [f(x, c) + f(x, d)] dx + \frac{1}{d - c} \int_c^d [f(a, y) + f(b, y)] dy \right].$$

*Proof.* From Lemma 1, we have

$$\begin{aligned}
& \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
& \quad \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\
& \leq \frac{(b-a)(d-c)}{4} \\
& \quad \times \int_0^1 \int_0^1 |(1-2t)(1-2s)| \left| \frac{\partial^2 f}{\partial t \partial s}(ta + (1-t)b, sc + (1-s)d) \right| dt ds.
\end{aligned}$$

Since  $f : \Delta \rightarrow \mathbb{R}$  is co-ordinated convex on  $\Delta$ , then one has:

$$\begin{aligned}
& \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
& \quad \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\
& \leq \frac{(b-a)(d-c)}{4} \\
& \quad \times \int_0^1 \left[ \int_0^1 |(1-2t)(1-2s)| \left\{ t \left| \frac{\partial^2 f}{\partial t \partial s}(a, sc + (1-s)d) \right| \right. \right. \\
& \quad \left. \left. + (1-t) \left| \frac{\partial^2 f}{\partial t \partial s}(b, sc + (1-s)d) \right| \right\} dt \right] ds.
\end{aligned}$$

Firstly, by calculating the integral in above inequality, we have

$$\begin{aligned}
& \int_0^1 |1-2t| \left\{ t \left| \frac{\partial^2 f}{\partial t \partial s}(a, sc + (1-s)d) \right| + (1-t) \left| \frac{\partial^2 f}{\partial t \partial s}(b, sc + (1-s)d) \right| \right\} dt \\
& = \int_0^{\frac{1}{2}} (1-2t) \left\{ t \left| \frac{\partial^2 f}{\partial t \partial s}(a, sc + (1-s)d) \right| + (1-t) \left| \frac{\partial^2 f}{\partial t \partial s}(b, sc + (1-s)d) \right| \right\} dt \\
& \quad + \int_{\frac{1}{2}}^1 (2t-1) \left\{ t \left| \frac{\partial^2 f}{\partial t \partial s}(a, sc + (1-s)d) \right| + (1-t) \left| \frac{\partial^2 f}{\partial t \partial s}(b, sc + (1-s)d) \right| \right\} dt \\
& = \frac{1}{4} \left( \left| \frac{\partial^2 f}{\partial t \partial s}(a, sc + (1-s)d) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(b, sc + (1-s)d) \right| \right).
\end{aligned}$$

Thus, we obtain

$$\begin{aligned}
 (2.5) \quad & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\
 & \leq \frac{(b-a)(d-c)}{16} \\
 & \times \int_0^1 |1-2s| \left\{ \left| \frac{\partial^2 f}{\partial t \partial s}(a, sc + (1-s)d) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(b, sc + (1-s)d) \right| \right\} ds.
 \end{aligned}$$

A similar way for other integral, since  $f : \Delta \rightarrow \mathbb{R}$  is co-ordinated convex on  $\Delta$ , we get

$$\begin{aligned}
 (2.6) \quad & \int_0^1 |1-2s| \left\{ \left| \frac{\partial^2 f}{\partial t \partial s}(a, sc + (1-s)d) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(b, sc + (1-s)d) \right| \right\} ds \\
 & = \int_0^{\frac{1}{2}} (1-2s) \left\{ s \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right| + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right| \right\} ds \\
 & \quad + \int_0^{\frac{1}{2}} (1-2s) \left\{ s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right| + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right| \right\} ds \\
 & = \int_{\frac{1}{2}}^1 (2s-1) \left\{ s \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right| + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right| \right\} ds \\
 & \quad + \int_{\frac{1}{2}}^1 (2s-1) \left\{ s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right| + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right| \right\} ds \\
 & = \frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|}{4}.
 \end{aligned}$$

By the (2.5) and (2.6), we get the inequality (2.4).  $\square$

**Theorem 3.** Let  $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a partial differentiable mapping on  $\Delta := [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b$  and  $c < d$ . If  $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q$ ,  $q > 1$ , is a convex function on

the co-ordinates on  $\Delta$ , then one has the inequalities:

$$\begin{aligned}
 (2.7) \quad & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\
 & \leq \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}} \\
 & \times \left( \frac{\left| \frac{\partial^2 f}{\partial s \partial t}(a, c) \right|^q + \left| \frac{\partial^2 f}{\partial s \partial t}(a, d) \right|^q + \left| \frac{\partial^2 f}{\partial s \partial t}(b, c) \right|^q + \left| \frac{\partial^2 f}{\partial s \partial t}(b, d) \right|^q}{4} \right)^{\frac{1}{q}}
 \end{aligned}$$

where

$$A = \frac{1}{2} \left[ \frac{1}{b-a} \int_a^b [f(x, c) + f(x, d)] dx + \frac{1}{d-c} \int_c^d [f(a, y) + f(b, y)] dy \right].$$

and  $\frac{1}{p} + \frac{1}{q} = 1$ .

*Proof.* From Lemma 1, we have

$$\begin{aligned}
 & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\
 & \leq \frac{(b-a)(d-c)}{4} \\
 & \times \int_0^1 \int_0^1 |(1-2t)(1-2s)| \left| \frac{\partial^2 f}{\partial t \partial s}(ta + (1-t)b, sc + (1-s)d) \right| dt ds.
 \end{aligned}$$

By using the well known Hölder inequality for double integrals,  $f : \Delta \rightarrow \mathbb{R}$  is co-ordinated convex on  $\Delta$ , then one has:

$$\begin{aligned}
 & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\
 & \leq \frac{(b-a)(d-c)}{4} \left( \int_0^1 \int_0^1 |(1-2t)(1-2s)|^p dt ds \right)^{\frac{1}{p}} \\
 & \times \left( \int_0^1 \int_0^1 \left| \frac{\partial^2 f}{\partial t \partial s}(ta + (1-t)b, sc + (1-s)d) \right|^q dt ds \right)^{\frac{1}{q}}.
 \end{aligned}$$

Since  $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q$  is convex function on the co-ordinates on  $\Delta$ , we know that for  $t \in [0, 1]$

$$\begin{aligned} & \left| \frac{\partial^2 f}{\partial t \partial s}(ta + (1-t)b, sc + (1-s)d) \right|^q \\ & \leq t \left| \frac{\partial^2 f}{\partial t \partial s}(a, sc + (1-s)d) \right|^q + (1-t) \left| \frac{\partial^2 f}{\partial t \partial s}(b, sc + (1-s)d) \right|^q \end{aligned}$$

and

$$\begin{aligned} & \left| \frac{\partial^2 f}{\partial t \partial s}(ta + (1-t)b, sc + (1-s)d) \right|^q \\ & \leq ts \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + t(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q \\ & \quad + (1-t)s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-t)(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \end{aligned}$$

hence, it follows that

$$\begin{aligned} & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\ & \quad \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\ & \leq \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}} \\ & \quad \times \left( \int_0^1 \int_0^1 \left\{ ts \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + t(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q \right. \right. \\ & \quad \left. \left. + (1-t)s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-t)(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \right\} dt ds \right)^{\frac{1}{q}} \\ & = \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}} \\ & \quad \times \left( \frac{\left| \frac{\partial^2 f}{\partial s \partial t}(a, c) \right|^q + \left| \frac{\partial^2 f}{\partial s \partial t}(a, d) \right|^q + \left| \frac{\partial^2 f}{\partial s \partial t}(b, c) \right|^q + \left| \frac{\partial^2 f}{\partial s \partial t}(b, d) \right|^q}{4} \right)^{\frac{1}{q}}. \end{aligned}$$

□

**Theorem 4.** Let  $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a partial differentiable mapping on  $\Delta := [a, b] \times [c, d]$  in  $\mathbb{R}^2$  with  $a < b$  and  $c < d$ . If  $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q$ ,  $q \geq 1$ , is a convex function on

the co-ordinates on  $\Delta$ , then one has the inequalities:

$$\begin{aligned}
 (2.8) \quad & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\
 & \leq \frac{(b-a)(d-c)}{16} \\
 & \times \left( \frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{4} \right)^{\frac{1}{q}}
 \end{aligned}$$

where

$$A = \frac{1}{2} \left[ \frac{1}{b-a} \int_a^b [f(x, c) + f(x, d)] dx + \frac{1}{d-c} \int_c^d [f(a, y) + f(b, y)] dy \right].$$

*Proof.* From Lemma 1, we have

$$\begin{aligned}
 & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\
 & \leq \frac{(b-a)(d-c)}{4} \\
 & \times \int_0^1 \int_0^1 |(1-2t)(1-2s)| \left| \frac{\partial^2 f}{\partial t \partial s}(ta + (1-t)b, sc + (1-s)d) \right| dt ds.
 \end{aligned}$$

By using the well known power mean inequality for double integrals,  $f : \Delta \rightarrow \mathbb{R}$  is co-ordinated convex on  $\Delta$ , then one has:

$$\begin{aligned}
 & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\
 & \leq \frac{(b-a)(d-c)}{4} \left( \int_0^1 \int_0^1 |(1-2t)(1-2s)| dt ds \right)^{1-\frac{1}{q}} \\
 & \times \left( \int_0^1 \int_0^1 |(1-2t)(1-2s)| \left| \frac{\partial^2 f}{\partial t \partial s}(ta + (1-t)b, sc + (1-s)d) \right|^q dt ds \right)^{\frac{1}{q}}.
 \end{aligned}$$

Since  $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q$  is convex function on the co-ordinates on  $\Delta$ , we know that for  $t \in [0, 1]$

$$\begin{aligned} & \left| \frac{\partial^2 f}{\partial t \partial s}(ta + (1-t)b, sc + (1-s)d) \right|^q \\ \leq & t \left| \frac{\partial^2 f}{\partial t \partial s}(a, sc + (1-s)d) \right|^q + (1-t) \left| \frac{\partial^2 f}{\partial t \partial s}(b, sc + (1-s)d) \right|^q \end{aligned}$$

and

$$\begin{aligned} & \left| \frac{\partial^2 f}{\partial t \partial s}(ta + (1-t)b, sc + (1-s)d) \right|^q \\ \leq & ts \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + t(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q \\ & + (1-t)s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-t)(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \end{aligned}$$

hence, it follows that

$$\begin{aligned} & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\ & \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\ \leq & \frac{(b-a)(d-c)}{4} \left( \frac{1}{4} \right)^{1-\frac{1}{q}} \\ & \times \left( \int_0^1 \int_0^1 |(1-2t)(1-2s)| \left\{ ts \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + t(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q \right. \right. \\ & \left. \left. + (1-t)s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-t)(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \right\} dt ds \right)^{\frac{1}{q}}. \end{aligned}$$

Firstly, by calculating the integral in above inequality, we have

$$\begin{aligned}
& \int_0^1 |1-2t| \left( ts \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + t(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q \right. \\
& \quad \left. + (1-t)s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-t)(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \right) dt \\
&= \int_0^{\frac{1}{2}} (1-2t) \left( ts \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + t(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q \right. \\
& \quad \left. + (1-t)s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-t)(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \right) dt \\
& \quad + \int_{\frac{1}{2}}^1 (2t-1) \left( ts \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + t(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q \right. \\
& \quad \left. + (1-t)s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-t)(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \right) dt \\
&= \frac{s \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q}{24} + \frac{(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q}{24} \\
& \quad + \frac{5s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q}{24} + \frac{5(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{24} \\
& \quad + \frac{5s \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q}{24} + \frac{5(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q}{24} \\
& \quad + \frac{s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q}{24} + \frac{(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{24} \\
&= \frac{s \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q}{4} \\
& \quad + \frac{s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{4}.
\end{aligned}$$

Thus, we obtain

$$\begin{aligned}
 (2.9) \quad & \left| \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4} \right. \\
 & \left. + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dy dx - A \right| \\
 \leq & \frac{(b-a)(d-c)}{16} \left[ \int_0^1 |1-2s| \left( s \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q \right. \right. \\
 & \left. \left. + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \right) ds \right]^{\frac{1}{q}}.
 \end{aligned}$$

A similar way for other integral, since  $f : \Delta \rightarrow \mathbb{R}$  is co-ordinated convex on  $\Delta$ , we get

$$\begin{aligned}
 (2.10) \quad & \int_0^1 |1-2s| \left( s \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q \right. \\
 & \left. + s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \right) ds \\
 = & \int_0^{\frac{1}{2}} (1-2s) \left( s \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q \right. \\
 & \left. + s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \right) ds \\
 & + \int_{\frac{1}{2}}^1 (2s-1) \left( s \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q \right. \\
 & \left. + s \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + (1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \right) ds \\
 = & \frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q}{24} + \frac{5 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q}{24} + \frac{\left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q}{24} + \frac{5 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{24} \\
 & + \frac{5 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q}{24} + \frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q}{24} + \frac{5 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q}{24} + \frac{\left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{24} \\
 = & \frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q}{4} + \frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q}{4} + \frac{\left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q}{4} + \frac{\left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{4}.
 \end{aligned}$$

By the (2.9) and (2.10), we get the inequality (2.8).  $\square$

**Remark 1.** Since  $\frac{1}{4} < \frac{1}{(p+1)^{\frac{2}{p}}} < 1$ , if  $p > 1$ , the estimation given in Theorem 4 is better than the one given in Theorem 3.

## REFERENCES

- [1] M. Alomari and M. Darus, Co-ordinated  $s$ -convex function in the first sense with some Hadamard-type inequalities, *Int. J. Contemp. Math. Sciences*, 3 (32) (2008), 1557-1567.
- [2] M. Alomari and M. Darus, On the Hadamard's inequality for  $\log$ -convex functions on the coordinates, *J. of Inequal. and Appl*, Article ID 283147, (2009), 13 pages.
- [3] S.S. Dragomir, On Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane, *Taiwanese Journal of Mathematics*, 4 (2001), 775-788.
- [4] M. A. Latif and M. Alomari, Hadamard-type inequalities for product two convex functions on the co-ordinates, *Int. Math. Forum*, 4(47), 2009, 2327-2338.
- [5] M. A. Latif and M. Alomari, On the Hadamard-type inequalities for  $h$ -convex functions on the co-ordinates, *Int. J. of Math. Analysis*, 3(33), 2009, 1645-1656.
- [6] M.E. Özdemir, E. Set and M.Z. Sarikaya, New some Hadamard's type inequalities for co-ordinated  $m$ -convex and  $(\alpha, m)$ -convex functions, *RGMI*A, Res. Rep. Coll., 13 (2010), Supplement, Article 4.

♣DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE, TURKEY

*E-mail address:* sarikayamz@gmail.com

■ATATÜRK UNIVERSITY, K.K. EDUCATION FACULTY, DEPARTMENT OF MATHEMATICS, 25240, CAMPUS, ERZURUM, TURKEY

*E-mail address:* erhanset@yahoo.com

♦GRADUATE SCHOOL OF NATURAL AND APPLIED SCIENCES, AĞRI İBRAHİM ÇEÇEN UNIVERSITY, AĞRI, TURKEY

*E-mail address:* emos@atauni.edu.tr

▼RESEARCH GROUP IN MATHEMATICAL INEQUALITIES & APPLICATIONS, SCHOOL OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO Box 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

*E-mail address:* sever.dragomir@vu.edu.au

*URL:* <http://rgmia.vu.edu.au/dragomir>